

PHB-degrading bacteria isolated from the gastrointestinal tract of aquatic animals as protective actors against luminescent vibriosis

Yiying Liu¹, Peter De Schryver², Bart Van Delsen¹, Loïs Maignien², Nico Boon², Patrick Sorgeloos¹, Willy Verstraete², Peter Bossier¹ & Tom Defoirdt^{1,2}

¹Laboratory of Aquaculture and Artemia Reference Center, Ghent University, Rozier, Ghent, Belgium; and ²Laboratory for Microbial Ecology and Technology (LabMET), Ghent University, Coupure Links, Ghent, Belgium

Correspondence: Peter Bossier, Laboratory of Aquaculture and Artemia Reference Center, Ghent University, Rozier 44, B-9000 Ghent, Belgium. Tel.: +32 9 264 37 54; fax: +32 9 264 41 93; e-mail: peter.bossier@UGent.be

Received 13 April 2010; revised 21 May 2010; accepted 21 May 2010.
Final version published online 28 June 2010.

DOI:10.1111/j.1574-6941.2010.00926.x

Editor: Julian Marchesi

Keywords

polyhydroxyalkanoate; PHB; depolymerization; brine shrimp; probiotic; aquaculture.

Abstract

The use of poly- β -hydroxybutyrate (PHB) was shown to be successful in increasing the resistance of brine shrimp against pathogenic infections. In this study, we isolated for the first time PHB-degrading bacteria from a gastrointestinal environment. Pure strains of PHB-degrading bacteria were isolated from Siberian sturgeon, European sea bass and giant river prawn. The capability of selected isolates to degrade PHB was confirmed in at least two of three setups: (1) growth in minimal medium containing PHB as the sole carbon (C) source, (2) production of clearing zones on minimal agar containing PHB as the sole C source and (3) degradation of PHB (as determined by HPLC analysis) in 10% Luria–Bertani medium containing PHB. Challenge tests showed that the PHB-degrading activity of the selected isolates increased the survival of brine shrimp larvae challenged to a pathogenic *Vibrio campbellii* strain by a factor 2–3. Finally, one of the PHB-degrading isolates from sturgeon showed a double biocontrol effect because it was also able to inactivate acylhomoserine lactones, a type of quorum-sensing molecule that regulates the virulence of different pathogenic bacteria. Thus, the combined supplementation of a PHB-degrading bacterium and PHB as a synbioticum provides perspectives for improving the gastrointestinal health of aquatic animals.

Introduction

The prophylactic use of antibiotics in aquaculture has resulted in multiple resistance in several pathogens, leading to ineffective treatment and an increased risk of resistance transfer to animal and human pathogens (Defoirdt *et al.*, 2007a; Das *et al.*, 2009). Also, trace amounts of antibiotics have been detected in aquatic products for human consumption and unwanted alterations of bacterial communities in sediment and water column have been observed (Cabello, 2006). Consequently, there is a growing awareness that antibiotics should be used more carefully in animal production and the prophylactic use should be stopped. However, a decreased use of antibiotics could result in a higher frequency of pathogenic bacteria, which in turn could lead to a higher incidence of infections.

Several research groups are currently investigating biological and environment-friendly alternative approaches to fight disease problems in aquaculture (Defoirdt *et al.*, 2007a).

In this respect, currently, there is considerable interest in short-chain fatty acids as biocontrol agents in animal production. However, there are some important limitations to the use of short-chain fatty acids (especially in aquaculture settings, where they can diffuse into the culture water). Therefore, we have recently started investigations on the use of polyhydroxyalkanoates, polymers of β -hydroxy short-chain fatty acids, as an alternative (for a review, see Defoirdt *et al.*, 2009). The bacterial storage compound poly- β -hydroxybutyrate (PHB) is the most well-known polyhydroxyalkanoate. This compound is accumulated as a carbon (C) and energy storage reserve by a wide variety of bacteria residing in a broad range of habitats (Muller & Seebach, 1993). The use of PHB as an anti-infective strategy has been shown in studies using gnotobiotic *Artemia franciscana* (Defoirdt *et al.*, 2007b; Halet *et al.*, 2007). PHB is not water soluble, and consequently, it needs to be degraded into water-soluble products (i.e. β -hydroxybutyrate monomers and oligomers) in the gastrointestinal tract in order to have a beneficial effect on

the host (Defoirdt *et al.*, 2009). One strategy to improve the degradation of PHB into water-soluble products is to use PHB-degrading microorganisms, which have been shown before to increase the protective effect of PHB particles (Defoirdt *et al.*, 2007b).

Aerobic and anaerobic PHA-degrading microorganisms have been isolated from various environments (Jendrossek & Handrick, 2002; Calabia & Tokiwa, 2004). However, to the best of our knowledge, no such microorganisms have been isolated from gastrointestinal environments thus far. In this study, we report the isolation of PHB-degrading strains from the gastrointestinal environment of aquatic animals. Siberian sturgeon, European sea bass and giant river prawn were selected due to their importance for aquaculture and because they live in different environments (fresh water, seawater and brackish water, respectively) and have different feeding habits. The possibility of applying the isolates as protective agents was investigated in a model system with gnotobiotic brine shrimp (*A. franciscana*) challenged to a virulent *Vibrio campbellii* strain, in which the beneficial effects of PHB and PHB-degrading bacteria have been documented before (Defoirdt *et al.*, 2007b).

Materials and methods

Origin of intestinal microbiota and enrichment of PHB degraders

Samples were taken from the posterior gut content of 5-month-old Siberian sturgeon (*Acipenser baerii*). The fish had been fed at 3% on wet body weight day⁻¹ for 45 days with an artificial diet (Joosen-Luyckx Aqua Bio, Turnhout, Belgium) containing 5% PHB (w/w). Secondly, fecal matter of 3-month-old European sea bass (*Dicentrarchus labrax*) juveniles was collected. These had been fed at 3% on wet body weight day⁻¹ for 40 days with an artificial diet (Skretting, Boxmeer, the Netherlands) containing 10% PHB (w/w). Finally, intestinal microbiota were collected from larvae of the giant river prawn (*Macrobrachium rosenbergii*). These larvae had been fed *ad libitum* for 15 days with *Artemia* nauplii enriched for 24 h in a 5 g L⁻¹ PHB solution. After sampling, all bacterial sources were stored in 20% glycerol at -80 °C.

Before use, the intestinal microbiota were washed three times in a salt solution [5.0 g L⁻¹ synthetic sea salt (Instant Ocean[®], Aquarium Systems Inc., Sarrebourg, France)]. Each washed sample was inoculated in 10 mL minimal medium containing PHB as the sole C source. The medium contained 5.0 g L⁻¹ synthetic sea salt, 0.2 g L⁻¹ NH₄Cl and 1.0 g L⁻¹ PHB particles (average diameter 25 µm; Goodfellow Cambridge Ltd, Huntingdon, UK). The cultures were incubated at 28 °C on an orbital shaker at 130 r.p.m. After 3 days, each of the cultures was transferred at a concentration

of *c.* 10⁵ CFU mL⁻¹ into 100 mL of a fresh medium. This procedure was repeated to enrich PHB-degrading bacteria.

Isolation of PHB-degrading strains

The PHB-degrading mixed enrichment cultures were plated on Luria-Bertani (LB) agar and the morphologically most abundant colonies were picked using a sterile inoculation loop. Each colony type was incubated in minimal medium containing PHB for 72 h and subsequently purified by streak plating on LB agar. In this way, 38 pure cultures were obtained. Based on preliminary tests (growth in minimal medium containing PHB and effect toward *Artemia*), the six most promising strains were retained for further experimenting: S4 and S7 from sturgeon, B7 and B12 from sea bass and M13 and M15 from giant river prawn.

Identification of the isolates by 16S rRNA gene sequencing

PCR targeting a 1500-bp fragment of the 16S rRNA gene of the isolates was performed according to Boon *et al.* (2002) using the primer pair GM3f and GM4r (Biolegio, Nijmegen, the Netherlands) (Muyzer *et al.*, 1995). PCR was performed using a GeneAmps PCR system 2700 thermal cycler (PE Applied Biosystems, Nieuwerkerken a/d IJssel, the Netherlands) using the following program: 95 °C for 5 min, 32 cycles of 94 °C for 1 min, 42 °C for 1 min, 72 °C for 3 min and finally an extension period of 72 °C for 10 min.

DNA sequencing of the PCR products obtained was carried out at IIT Biotech (Bielefeld, Germany). The forward and reverse sequences obtained with the sequencing primers GM3f and GM4r were processed using EBIOX software, and homology searches were completed with the BLAST server of the National Centre for Biotechnology Information using the BLAST algorithm (Altschul *et al.*, 1997) for the comparison of the nucleotide query sequence against a nucleotide sequence database (BLASTN). The nucleotide sequences of the isolates were deposited in the GenBank database (<http://www.ncbi.nlm.nih.gov/Genbank>) with the accession numbers GU372408, GU372409, GU372410, GU372411, GU372412 and GU372413 for the strains S4, S7, B7, B12, M13 and M15, respectively.

Extracellular PHB depolymerase assay

Extracellular PHB depolymerase production was assessed qualitatively by spotting strains on solid medium containing PHB particles as the sole C source as described previously (Defoirdt *et al.*, 2007b). The medium contained 500 mg L⁻¹ PHB particles, 1 g L⁻¹ NH₄Cl, 5 g L⁻¹ artificial sea salt and 15 g L⁻¹ agar. The plates were incubated at 28 °C and examined daily for the presence of a clearing zone around the colonies. *Comamonas testosteroni* LMG 19554 was used as a positive control (Defoirdt *et al.*, 2007b).

Degradation of PHB in a nutrient-rich background

In order to study whether the PHB-degrading bacteria could degrade PHB in a nutrient-rich background, the bacteria were inoculated in 10% LB medium containing 1.25 g L^{-1} PHB at an initial density of 10^5 CFU mL^{-1} . After 3 days of incubation on a shaker at 28°C , residual PHB concentrations were determined by HPLC as described by De Schryver & Verstraete (2009). Briefly, samples were thawed in warm water and centrifuged for 10 min at 7000 g . The pellets were resuspended in 1 mL of distilled water and transferred to Eppendorf tubes. Samples were centrifuged for 5 min at $13\,000 \text{ g}$ and dried overnight at 100°C . Dried pellets were digested with 1 mL of 96% H_2SO_4 at 100°C for 1 h to form crotonic acid. The reaction mixture was then cooled to room temperature on ice, diluted 50-fold with $0.014 \text{ M H}_2\text{SO}_4$ and crotonic acid concentrations were determined by HPLC using a Dionex ASI-100 autosampler injector (Dionex Corporation, Sunnyvale, CA) equipped with an Aminex HPX-87 H ion-exchange organic acids column ($300 \times 7.8 \text{ mm}$) (Bio-Rad Laboratories, Richmond, CA). The solvent used was $0.014 \text{ M H}_2\text{SO}_4$ at a flow rate of 0.7 mL min^{-1} . The elution peaks were monitored at 210 nm using a Dionex UV detector. The PHB content was calculated from a calibration curve for standards of commercial PHB (Goodfellow Cambridge Ltd) treated in the same manner as the samples.

N-hexanoyl-L-homoserine lactone (HHL) inactivation assay

Autoclaved LB broth was supplemented with a $0.2\text{-}\mu\text{m}$ filter-sterilized HHL (Sigma-Aldrich, Germany) solution to a final concentration of 10 mg HHL L^{-1} [buffered at pH 6.5 by the addition of 3-(*N*-morpholino)-propanesulfonic acid]. Each strain was inoculated at 10^6 CFU mL^{-1} and incubated on an orbital shaker (130 r.p.m.) at 28°C . After 48 h, the culture was $0.2\text{-}\mu\text{m}$ filter-sterilized and the remaining HHL was measured by an agar diffusion assay using *Chromobacterium violaceum* CV026 as the reporter strain as described before (Baruah *et al.*, 2009). *Pseudomonas* spp. P3/pME6000 and *Pseudomonas* spp. P3/pME6863 (= pME6000 + the *Bacillus* AHL lactonase gene *aiiA*), grown under the same culture conditions, were used as the negative and positive control, respectively (Dang *et al.*, 2009).

Culture preparation for *Artemia in vivo* challenge tests

Isolates B7, B12, M13 and M15 were inoculated in marine broth (Difco Laboratories, Detroit), and S4 and S7 in LB broth. After incubation for 24 h at 28°C on an orbital shaker at 130 r.p.m., the cultures were washed with autoclaved

synthetic seawater and added to the *Artemia* culture at $c. 10^7 \text{ CFU mL}^{-1}$.

Artemia in vivo challenge tests

Artemia franciscana (EG[®] Type, INVE Aquaculture, Belgium) nauplii were hatched axenically as described by Defoirdt *et al.* (2005). After hatching, groups of 20 nauplii were transferred into sterile 50-mL cylindrical tubes containing 20 mL filtered and autoclaved artificial seawater. The animals were challenged with 10^5 CFU mL^{-1} *V. campbellii* LMG 21363, added to the *Artemia* culture water. At the start of the challenge test, a suspension of autoclaved LVS3 bacteria (Verschuere *et al.*, 1999) was added as feed at a concentration of 10^7 CFU mL^{-1} to the culture water. In a first group of challenge tests, challenged nauplii were either untreated or treated with the isolates (10^7 CFU mL^{-1}), with and without PHB (100 mg L^{-1}). In a second challenge test, challenged nauplii were treated with PHB (100 mg L^{-1}) in combination with either live or dead isolates. After addition of feed, PHB (100 mg L^{-1}), isolates and pathogen, tubes were placed on a rotor at 4 r.p.m. under continuous light at 28°C . After 48 h, the survival was determined. All treatments were performed in triplicate.

Statistics

Statistical analyses were carried out using the SPSS statistical software (version 15). Survival data of *Artemia* were compared using one-way ANOVA, followed by Duncan's *post hoc* test. For all statistical analyses, a 5% significance level was used.

Results

Enrichment of PHB-degrading bacteria from intestinal microbiota

Intestinal microbiota were collected from sturgeon, sea bass and giant river prawn that had received PHB in their diets. The microbiota were enriched for PHB degraders in a medium containing PHB as the sole C source. In this medium, cell concentrations rapidly increased with 2–3 log units within 24 h after inoculation (Fig. 1a). This was similar to the growth of *C. testosteroni* LMG 19554, which served as a positive control (Defoirdt *et al.*, 2007b). This indicated that the enrichment cultures were able to (partially) degrade the PHB particles into water-soluble products.

In order to confirm PHB-degrading activity, the enrichment cultures were spotted on minimal agar containing PHB particles as the sole C source. After incubation at 28°C , clearing zones were observed for the sturgeon and sea bass microbiota, but not for the prawn microbiota (Fig. 2a–c). The production of clearing zones requires complete degradation of the PHB particles, and the absence of clearing




Fig. 1. Growth of intestinal microbiota (a) and selected isolates (b) in minimal medium containing PHB as the sole C source. Noninoculated medium was used as a control. The data points represent the mean values of three replicates. Error bars are too small to be visible.

zones for the prawn microbiota suggests weaker PHB-degrading activity under these conditions. In a further experiment, we studied the degradation of PHB in a nutrient-rich background (10% LB medium). The addition of 10% LB medium simulated the presence of other easily degradable C sources in addition to PHB, as will be the case in the intestinal environment of aquaculture animals that contains feed. After 3 days of incubation at 28 °C on an orbital shaker, a very low residual PHB concentration was measured in the cultures of the sturgeon microbiota (Table 1). Intermediate concentrations were measured in cultures of the sea bass microbiota and the highest ones in those of the prawn microbiota, which is consistent with the weaker PHB-degrading activity of these microbiota as observed on the plates with PHB as the sole C source.

Isolation of PHB-degrading strains from the enrichment cultures

The PHB-degrading enrichment cultures were spread-plated on LB agar; dominant colonies were picked and further purified. Based on preliminary tests, six isolates were retained

for further experiments: S4 and S7 from sturgeon, B7 and B12 from sea bass and M13 and M15 from giant river prawn. The isolates B7, B12, M13 and M15 grew rapidly in minimal medium containing PHB as the sole C source, with a 2–3 log units increase in cell density after 24 h of incubation (Fig. 1b). It was not possible to accurately measure the cell density for isolates S4 and S7 because they were growing as flocs. However, these isolates showed clearing zone production on minimal agar containing PHB particles as the sole C source (Fig. 2d and e), indicating that they were able to degrade PHB.

The ability of the isolates to degrade PHB in a nutrient-rich background (10% LB medium) was investigated. After 3 days of incubation at 28 °C, the residual PHB concentrations were clearly lower than that in the negative (noninoculated) control for all isolates (Table 1).

Identification of the isolates

The six selected isolates obtained from the enrichment cultures were grown in LB medium, and a 1500-bp fragment of the 16S rRNA gene was amplified by PCR and sequenced. Homology searches with the BLAST server of the National Centre for Biotechnology Information showed that the isolates originating from the same host species were closely related (97.1%, 99.1% and 98.9% identity between the two strains isolated from sturgeon, sea bass and prawn, respectively). Isolates S4 and S7 (from sturgeon) were closely related to *Acidovorax* spp. (99.0% and 96.9% identity, respectively); B7 and B12 (from sea bass) were closely related to *Acinetobacter* spp. (99.4% and 99.2% identity, respectively); and M13 and M15 (from prawn) were closely related to *Ochrobactrum* spp. (99.5% and 99.3% identity, respectively).

Acylhomoserine lactone (AHL) inactivation assay

AHL degradation by the selected isolates was screened by growing them in LB medium containing 10 mg L⁻¹ HHL. After 48 h of incubation, the remaining HHL in filter-sterilized supernatants was measured in an agar diffusion assay with the AHL reporter strain *C. violaceum* CV026. In this assay, the presence of AHLs in the supernatant is manifested by purple pigment production by the reporter strain. The supernatants of isolate S7 did not induce pigment production in the reporter, indicating complete degradation of HHL (Fig. 3). For the other isolates, a purple zone with approximately the same diameter as the non-HHL-degrading control occurred, indicating no or very little HHL inactivation.

Effect of the isolates on the survival of brine shrimp nauplii challenged to pathogenic *V. campbellii*

In order to obtain a first indication of the potential of the selected isolates to protect cultured organisms from




Fig. 2. PHB degradation on minimal medium containing PHB particles as the sole C source. Complete degradation of the PHB particles results in clearing of the medium. (a) Intestinal microbiota from sturgeon, (b) intestinal microbiota from sea bass, (c) intestinal microbiota from prawn, (d) isolate S4, (e) isolate S7 and (f) the known PHB-depolymerizing bacterium *Comamonas testosteroni* LMG 19554.

Table 1. Residual PHB concentration after culturing intestinal microbiota and selected isolates for 72 h in 10% LB medium supplemented with 1.25 g L^{-1} PHB

| Inoculum | [PHB] after 72 h (g L^{-1}) |
|--|--|
| Negative control | 1.01 |
| Intestinal microbiota from sturgeon | 0.01 |
| Intestinal microbiota from sea bass | 0.63 |
| Intestinal microbiota from giant river prawn | 0.81 |
| Isolate S4 | 0.70 |
| Isolate S7 | 0.68 |
| Isolate B7 | 0.55 |
| Isolate B12 | 0.82 |
| Isolate M13 | 0.33 |
| Isolate M15 | 0.82 |

The initial microbial density was 10^5 CFU mL^{-1} . Uninoculated medium was used as a negative control.

infection, we performed challenge tests in a model system with gnotobiotic brine shrimp (*Artemia*) nauplii. In a first series of challenge tests, the protective effect of PHB degradation by the isolates against infection with pathogenic *V. campbellii* was investigated. The PHB-degrading isolates were added to the culture water either with or without 100 mg L^{-1} PHB. This PHB concentration was chosen because previous research at our laboratories showed that it




Fig. 3. HHL degradation by the isolates. Ten microliters filter-sterilized supernatants of cultures grown for 48 h in LB medium supplemented with 10 mg L^{-1} HHL were spotted on LB agar covered with a lawn of the AHL reporter strain *Chromobacterium violaceum* CV026. The presence of HHL in the supernatants is manifested by purple pigment production in the reporter strain. (a) Negative control (P3/pME6000), (b) positive control (P3/pME6863), (c) isolate S4 and (d) isolate S7.

was suboptimal in protecting *Artemia* against infection and that the addition of a PHB-degrading bacterium could enhance the protective effect of PHB at this concentration (Defoirdt *et al.*, 2007b). When added in combination with PHB, isolates B7, B12, M13 and M15 significantly increased the survival of *Artemia* nauplii when compared with untreated challenged nauplii (Fig. 4). For isolates S4 and S7, the survival was not significantly different. Therefore, these strains were not used in further challenge tests. The addition of the isolates without PHB did not result in a significant difference in survival compared with the challenged control without PHB, except for isolate B12.




Fig. 4. Percentage survival of *Artemia* nauplii after a 48-h challenge with *Vibrio campbellii* LMG 21363 (mean \pm SE of three replicates). The three different panels represent different challenge tests (performed at different times). The challenged nauplii were either untreated (control) or treated with a PHB-degrading isolate, with or without 100 mg L⁻¹ PHB. Within each panel, treatments with a different letter are significantly different ($P < 0.05$).

In a last challenge test, in order to confirm that the increased survival upon addition of the isolates was caused by PHB degradation and not by nutritional properties of the isolates, we studied the effect of adding the isolates dead to the culture water vs. adding them alive. The addition of live isolates resulted in a significantly higher survival of the *Artemia* when compared with the addition of the corresponding dead isolates, except for isolate B7 (Fig. 5). All the living isolates resulted in a higher survival than the challenged control, although the differences were not always significant.

Discussion

PHB has been shown to increase pathogen resistance and to beneficially influence the growth of aquatic animals in a number of studies (Defoirdt *et al.*, 2007b; Halet *et al.*, 2007; De Schryver *et al.*, 2009; Nhan *et al.*, 2010). A method to increase these beneficial effects could be the addition of extracellular PHB-degrading bacteria (Defoirdt *et al.*, 2009). PHB-degrading bacteria have been isolated from various environments such as soil, fresh and marine water, estuarine sediment and air (Jendrossek & Handrick, 2002). Bacteria that are used to beneficially affect a certain host need to persist and function in the specific ecological niche of the host gastrointestinal tract and, consequently, it is advisable to isolate them from this environment (or from a similar niche) in order to increase the chance of success for colonization (Reid *et al.*, 2003). Here, we isolated PHB-degrading bacteria from intestinal microbiota of Siberian sturgeon, European sea bass and giant river prawn that had received PHB in their diets. To our knowledge, this was the first attempt to isolate PHB-degrading strains from the intestinal tract of higher animals.

PHB-degrading pure strains were isolated from PHB-degrading enrichment cultures from the intestinal microbiota of the animals and their capability to degrade PHB was




Fig. 5. Percentage survival of *Artemia* nauplii after a 48-h challenge with *Vibrio campbellii* LMG 21363 (mean \pm SE of three replicates). The challenged nauplii were either not treated (control) or treated with living or dead isolates, in combination with 100 mg L⁻¹ PHB. Treatments with a different letter are significantly different ($P < 0.05$).

confirmed in at least two of three setups: (1) growth in minimal medium containing PHB as the sole C source, (2) production of clearing zones on minimal agar containing PHB as the sole C source and (3) degradation of PHB (as determined by HPLC analysis) in 10% LB medium containing PHB. The isolates originating from sea bass and giant river prawn grew well in the minimal medium containing PHB. The sturgeon isolates were not tested in this setup because they are growing as flocs in liquid medium (and thus growth could not be measured accurately by plating). However, their PHB-degrading capability was confirmed by the production of clearing zones on agar plates containing PHB particles. All isolates were able to degrade PHB in the last setup (although to a different extent), which simulated the presence of other, easily degradable, C sources as is the case in the gastrointestinal tract of animals. Interestingly, the isolates from sturgeon showed a lower PHB-degrading capacity than the mixed bacterial community from sea bass, suggesting that the cooperation between different strains might enhance the degradation rate. The PHB-degrading isolates originating from sturgeon belonged to the genus *Acidovorax*. Extracellular PHB-depolymerase-producing members of this genus have been isolated previously from pond water, river water and soil (Kobayashi *et al.*, 1999). The isolates from sea bass and giant river prawn belonged to the genus *Acinetobacter* and *Ochrobactrum*, respectively. Both genera have not been documented before to contain PHB-degrading species.

Previous research at our laboratories has shown that PHB can protect brine shrimp (*A. franciscana*) larvae from infection caused by the virulent pathogen *V. campbellii* LMG 21363 (Defoirdt *et al.*, 2007b; Halet *et al.*, 2007). However, in order to exert its beneficial effect, the polymer needs to be degraded into water-soluble products. The main end product of microbial PHB degradation is β -hydroxybutyrate (Jendrossek & Handrick, 2002), which has been reasoned to protect aquaculture animals from bacterial infections in two ways: (1) by providing energy to the intestinal mucosa, thereby increasing intestinal health and resistance to infections, and (2) by decreasing the growth and/or the virulence of the pathogens (Defoirdt *et al.*, 2009). The results presented in this study showed that the PHB-degrading activity of the selected isolates increased the survival of brine shrimp larvae challenged to the pathogenic *V. campbellii* strain. Indeed, the addition of the isolates together with 100 mg L^{-1} PHB generally resulted in a higher survival than the addition of the isolates without PHB, and the addition of live isolates with 100 mg L^{-1} PHB resulted in a higher survival than the addition of the corresponding dead isolates with 100 mg L^{-1} PHB. Only strains S4 and S7 (isolated from sturgeon) were not able to protect the brine shrimp larvae from the pathogen, which can be ascribed to two factors: (1) both strains were growing as flocs in liquid

medium and most of them were too large to be ingested by the brine shrimp larvae. Indeed, brine shrimp graze on particles with a diameter preferentially smaller than $50 \mu\text{m}$ (Sorgeloos *et al.*, 1986) and (2) these strains were isolated from a freshwater environment and were probably not active in the brine shrimp cultures (which were performed in seawater). Indeed, these isolates did not grow in LB medium with a high salt concentration (35 g L^{-1}). In contrast, the isolates from sea bass and giant river prawn were euryhaline (able to grow between 5 and 35 g L^{-1}), which might make them useful under a broad range of environmental conditions.

Quorum-sensing, bacterial cell-to-cell communication with small signal molecules has been shown to regulate the virulence of several pathogenic bacteria, including aquaculture pathogens (Defoirdt *et al.*, 2004). Several types of signal molecules have been identified, with the best-studied ones being AHLs. Previous research at our laboratories has shown that AHL-degrading bacteria can readily be isolated from the intestinal microbiota of aquatic animals and that these bacteria can protect cultured animals from bacterial infections (Dang *et al.*, 2009). Therefore, AHL degradation by the selected isolates was screened by growing them in LB medium containing 10 mg L^{-1} HHL. This experiment revealed that isolate S7 was able to degrade HHL. No AHL-degrading bacteria belonging to the genus *Acidovorax* have been reported thus far. However, the 16S rRNA gene of isolate S7 showed 92.6% identity with the sequence of *Variovorax paradoxus* S110 (GenBank accession number CP001635.1) and *V. paradoxus* is a known AHL-degrading species (Leadbetter & Greenberg, 2000). Hence, strain S7 displays two beneficial anti-infective characteristics (PHB degradation and quorum-sensing disruption), and consequently, although it did not reveal any positive effect toward brine shrimp larvae in this study, it certainly is worthwhile to evaluate its beneficial potential in its original host.

To conclude, this research was the first to report the isolation of PHB-degrading bacteria from a gastrointestinal environment. The addition of selected PHB-degrading isolates together with PHB significantly increased the survival of brine shrimp larvae infected with the virulent pathogen *V. campbellii* LMG 21363. The effects noted in the challenge tests represent an increased survival by a factor 2–3, which is highly relevant for practice. Moreover, the dose of the PHB involved (100 mg L^{-1}) is economically feasible. Finally, the isolates involved are not related to aquaculture disease-associated genera and are easy to propagate. Hence, they offer possibilities for production at a technically relevant scale. Detailed studies on the use of the strains as probiotic feed components for the animals from which they were isolated will be the subject of our further research. These studies will reveal whether the isolates are active in different hosts and whether they can establish and maintain

themselves in the intestinal tract of these hosts or whether repeated addition is necessary.

Acknowledgements

This work was performed and funded within the frame of the Research Foundation of Flanders (FWO) project 'Probiotic-induced functional responses in aquatic organisms' and the European FP7 project 'Promicrobe – Microbes as positive actors for more sustainable aquaculture' (project reference: 227197). Additional financial support was obtained from a Master Grant from Ghent University to Y.L. and an FWO Postdoc Grant to T.D.

Authors' contribution

Y.L. and P.D.S. contributed equally to this work.

References

- Altschul SF, Madden TL, Schaffer AA, Zhang JH, Zhang Z, Miller W & Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**: 3389–3402.
- Baruah K, Cam DTV, Dierckens K, Wille M, Defoirdt T, Sorgeloos P & Bossier P (2009) *In vivo* effects of single or combined *N*-acyl homoserine lactone quorum sensing signals on the performance of *Macrobrachium rosenbergii* larvae. *Aquaculture* **288**: 233–238.
- Boon N, De Windt W, Verstraete W & Top EM (2002) Evaluation of nested PCR-DGGE (denaturing gradient gel electrophoresis) with group-specific 16S rRNA primers for the analysis of bacterial communities from different wastewater treatment plants. *FEMS Microbiol Ecol* **39**: 101–112.
- Cabello FC (2006) Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environ Microbiol* **8**: 1137–1144.
- Calabia BP & Tokiwa Y (2004) Microbial degradation of poly (D -3-hydroxybutyrate) by a new thermophilic *Streptomyces* isolate. *Biotechnol Lett* **26**: 15–19.
- Dang TVC, Nguyen VH, Dierckens K, Defoirdt T, Boon N, Sorgeloos P & Bossier P (2009) Novel approach of using homoserine lactone-degrading and poly-beta-hydroxybutyrate-accumulating bacteria to protect *Artemia* from the pathogenic effects of *Vibrio harveyi*. *Aquaculture* **291**: 23–30.
- Das A, Saha D & Pal J (2009) Antimicrobial resistance and *in vitro* gene transfer in bacteria isolated from the ulcers of EUS-affected fish in India. *Lett Appl Microbiol* **49**: 497–502.
- Defoirdt T, Boon N, Bossier P & Verstraete W (2004) Disruption of bacterial quorum sensing: an unexplored strategy to fight infections in aquaculture. *Aquaculture* **240**: 69–88.
- Defoirdt T, Bossier P, Sorgeloos P & Verstraete W (2005) The impact of mutations in the quorum sensing systems of *Aeromonas hydrophila*, *Vibrio anguillarum* and *Vibrio harveyi* on their virulence towards gnotobiotically cultured *Artemia franciscana*. *Environ Microbiol* **7**: 1239–1247.
- Defoirdt T, Boon N, Sorgeloos P, Verstraete W & Bossier P (2007a) Alternatives to antibiotics to control bacterial infections: luminescent vibriosis in aquaculture as an example. *Trends Biotechnol* **25**: 472–479.
- Defoirdt T, Halet D, Vervaeren H, Boon N, Van de Wiele T, Sorgeloos P, Bossier P & Verstraete W (2007b) The bacterial storage compound poly-beta-hydroxybutyrate protects *Artemia franciscana* from pathogenic *Vibrio campbellii*. *Environ Microbiol* **9**: 445–452.
- Defoirdt T, Boon N & Bossier P (2009) Short-chain fatty acids and poly- β -hydroxyalkanoates: (New) biocontrol agents for a sustainable animal production. *Biotechnol Adv* **27**: 680–685.
- De Schryver P & Verstraete W (2009) Nitrogen removal from aquaculture pond water by heterotrophic nitrogen assimilation in lab-scale sequencing batch reactors. *Bioresource Technol* **100**: 1162–1167.
- De Schryver P, Sinha AK, Kunwar PS, Baruah K, Boon N, Verstraete W, De Boeck G & Bossier P (2009) Poly-beta-hydroxybutyrate (PHB) increases growth performance and intestinal bacterial range-weighted richness in juvenile European sea bass, *Dicentrarchus labrax*. *Appl Microbiol Biot* **86**: 1535–1541.
- Halet D, Defoirdt T, Van Damme P, Vervaeren H, Forrez I, Van de Wiele T, Boon N, Sorgeloos P, Bossier P & Verstraete W (2007) Poly-beta-hydroxybutyrate-accumulating bacteria protect gnotobiotic *Artemia franciscana* from pathogenic *Vibrio campbellii*. *FEMS Microbiol Ecol* **60**: 363–369.
- Jendrossek D & Handrick R (2002) Microbial degradation of polyhydroxyalkanoates. *Annu Rev Microbiol* **56**: 403–432.
- Kobayashi T, Sugiyama A, Kawase Y, Saito T, Mergaert J & Swings J (1999) Biochemical and genetic characterization of an extracellular poly(3-hydroxybutyrate) depolymerase from *Acidovorax* sp. strain TP4. *J Environ Polym Degr* **7**: 9–18.
- Leadbetter JR & Greenberg EP (2000) Metabolism of acyl-homoserine lactone quorum-sensing signals by *Variovorax paradoxus*. *J Bacteriol* **182**: 6921–6926.
- Muller HM & Seebach D (1993) Poly(hydroxyalkanoates) – a 5th class of physiologically important organic biopolymers. *Angew Chem Int Edit* **32**: 477–502.
- Muyzer G, Teske A, Wirsén CO & Jannasch HW (1995) Phylogenetic relationships of *Thiomicrospira* species and their identification in deep-sea hydrothermal vent samples by denaturing gradient gel-electrophoresis of 16S rDNA fragments. *Arch Microbiol* **164**: 165–172.
- Nhan DT, Wille M, De Schryver P, Defoirdt T, Bossier P & Sorgeloos P (2010) The effect of poly- β -hydroxybutyrate on larviculture of the giant freshwater prawn (*Macrobrachium rosenbergii*). *Aquaculture* **302**: 76–81.
- Reid G, Sanders ME, Gaskins HR, Gibson GR, Mercenier A, Rastall R, Roberfroid M, Rowland I, Cherbut C &

- Klaenhammer TR (2003) New scientific paradigms for probiotics and prebiotics. *J Clin Gastroenterol* **37**: 105–118.
- Sorgeloos P, Lavens P, Léger P, Tackaert W & Versichele D (1986) *Manual for the Culture and Use of Brine Shrimp Artemia in Aquaculture*. Artemia Reference Center, Ghent, Belgium.
- Verschuere L, Rombaut G, Huys G, Dhont J, Sorgeloos P & Verstraete W (1999) Microbial control of the culture of *Artemia* juveniles through preemptive colonization by selected bacterial strains. *Appl Environ Microb* **65**: 2527–2533.